

**Class XII Session 2024-25**  
**Subject - Mathematics**  
**Sample Question Paper - 4**

**Time Allowed: 3 hours**

**Maximum Marks: 80**

**General Instructions:**

1. This Question paper contains - five sections A, B, C, D and E. Each section is compulsory. However, there are internal choices in some questions.
2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

**Section A**

1. Let  $A = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$ , then  $A^n$  is equal to [1]

a)  $\begin{bmatrix} a^n & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$       b)  $\begin{bmatrix} na & 0 & 0 \\ 0 & na & 0 \\ 0 & 0 & na \end{bmatrix}$   
 c)  $\begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a^n \end{bmatrix}$       d)  $\begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a \end{bmatrix}$

2. If the matrix  $A = \begin{bmatrix} 3-2x & x+1 \\ 2 & 4 \end{bmatrix}$  is singular then  $x = ?$  [1]

a) 1      b) 0  
 c) -1      d) -2

3. If A and B are invertible matrices, then which of the following is not correct? [1]

a)  $(AB)^{-1} = B^{-1} A^{-1}$       b)  $(A + B)^{-1} = B^{-1} + A^{-1}$   
 c)  $\det(A)^{-1} = [\det(A)]^{-1}$       d)  $\text{adj } A = |A| \cdot A^{-1}$

4. Let  $f(x) = [x]^2 + \sqrt{x}$ , where  $[\bullet]$  and  $\{\bullet\}$  respectively denotes the greatest integer and fractional part functions, then [1]

a)  $f(x)$  is continuous and differentiable at  $x = 0$       b)  $f(x)$  is non differentiable  $\forall x \in \mathbb{Z}$   
 c)  $f(x)$  is discontinuous  $\forall x \in \mathbb{Z} - \{1\}$       d)  $f(x)$  is continuous at all integral points

5. Find the equation of the line which passes through the point  $(1, 2, 3)$  and is parallel to the vector  $3\hat{i} + 2\hat{j} - 2\hat{k}$ . [1]

a)  $\vec{r} = \hat{i} + 2\hat{j} + 3\hat{k} + \lambda (3\hat{i} + 2\hat{j} - 2\hat{k})$ ,  $\lambda \in R$

b)  $\vec{r} = \hat{2i} + 2\hat{j} + 3\hat{k} + \lambda (3\hat{i} + 2\hat{j} - 2\hat{k})$ ,  $\lambda \in R$

c)  $\vec{r} = 4\hat{i} + 2\hat{j} + 3\hat{k} + \lambda (3\hat{i} + 2\hat{j} - 2\hat{k})$ ,  $\lambda \in R$

d)  $\vec{r} = 3\hat{i} + 2\hat{j} + 3\hat{k} + \lambda (3\hat{i} + 2\hat{j} - 2\hat{k})$ ,  $\lambda \in R$

6. The order of the differential equation of all circles of given radius a is: [1]

a) 4 b) 1

c) 2 d) 3

7. By graphical method solution of LLP maximize  $Z = x + y$  subject to  $x + y \leq 2x; y \geq 0$  obtained at [1]

a) at infinite number of points b) only two points

c) only one point d) at definite number of points

8. The domain of the function  $\cos^{-1}(2x - 1)$  is [1]

a)  $[0, \pi]$  b)  $[-1, 1]$

c)  $[0, 1]$  d)  $(-1, 0)$

9.  $\int_0^{\pi/2} \frac{\cos x}{(2+\sin x)(1+\sin x)} dx$  equals [1]

a)  $\log\left(\frac{3}{4}\right)$  b)  $\log\left(\frac{3}{2}\right)$

c)  $\log\left(\frac{4}{3}\right)$  d)  $\log\left(\frac{2}{3}\right)$

10. If  $\begin{bmatrix} x & -5 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ 1 \end{bmatrix} = 0$ , then the value of x is [1]

a)  $\pm 6\sqrt{5}$  b)  $5\sqrt{5}$

c)  $\pm 4\sqrt{3}$  d)  $\pm 3\sqrt{5}$

11. Objective function of an LPP is [1]

a) a function to be optimized b) a function between the variables

c) a constraint d) a relation between the variables

12. The vector in the direction of the vector  $\hat{i} - 2\hat{j} + 2\hat{k}$  that has magnitude 9 is [1]

a)  $\hat{i} - 2\hat{j} + 2\hat{k}$  b)  $3(\hat{i} - 2\hat{j} + 2\hat{k})$

c)  $9(\hat{i} - 2\hat{j} + 2\hat{k})$  d)  $\frac{\hat{i} - 2\hat{j} + 2\hat{k}}{3}$

13. If  $A = \begin{vmatrix} 1 & 0 & 0 \\ 1 & 1 & 2 \\ 3 & -1 & 9 \end{vmatrix}$ , then the value of  $\det(\text{Adj}(\text{Adj } A))$  equals [1]

a) 14641 b) 121

c) 11 d) 1331

14. If A and B are independent events such that  $P(A) = \frac{1}{5}$ ,  $P(A \cup B) = \frac{7}{10}$ , then what is  $P(\bar{B})$  equal to? [1]

a)  $\frac{3}{8}$  b)  $\frac{7}{9}$

c)  $\frac{3}{7}$  d)  $\frac{2}{7}$

15. Degree of the differential equation  $\sin x + \cos\left(\frac{dy}{dx}\right) = y^2$  is [1]

a) 2 b) not defined

c) 0 d) 1

16. If  $|\vec{a} \times \vec{b}| = 4$ ,  $|\vec{a} \cdot \vec{b}| = 2$ , then  $|\vec{a}|^2 |\vec{b}|^2 =$  [1]

a) 2 b) 20

c) 8 d) 6

17. If  $y = \tan^{-1} \frac{\cos x}{1 + \sin x}$  then  $\frac{dy}{dx} = ?$  [1]

a)  $\frac{1}{2}$  b) 1

c) 0 d)  $\frac{-1}{2}$

18. The cartesian equation of a line is given by  $\frac{2x-1}{\sqrt{3}} = \frac{y+2}{2} = \frac{z-3}{3}$  [1]

The direction cosines of the line is

a)  $\frac{\sqrt{3}}{\sqrt{55}}, \frac{-4}{\sqrt{55}}, \frac{6}{\sqrt{55}}$  b)  $\frac{3}{\sqrt{55}}, \frac{4}{\sqrt{55}}, \frac{6}{\sqrt{55}}$

c)  $\frac{\sqrt{3}}{\sqrt{55}}, \frac{4}{\sqrt{55}}, \frac{6}{\sqrt{55}}$  d)  $\frac{-3}{\sqrt{55}}, \frac{4}{\sqrt{55}}, \frac{6}{\sqrt{55}}$

19. **Assertion (A):** If manufacturer can sell x items at a price of  $\text{₹}(5 - \frac{x}{100})$  each. The cost price of x items is  $\text{₹}(\frac{x}{5} + 500)$ . Then, the number of items he should sell to earn maximum profit is 240 items. [1]

**Reason (R):** The profit for selling x items is given by  $\frac{24}{5}x - \frac{x^2}{100} - 300$ .

a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.

c) A is true but R is false. d) A is false but R is true.

20. **Assertion (A):** Let  $A = \{1, 5, 8, 9\}$ ,  $B = \{4, 6\}$  and  $f = \{(1, 4), (5, 6), (8, 4), (9, 6)\}$ , then f is a bijective function. [1]

**Reason (R):** Let  $A = \{1, 5, 8, 9\}$ ,  $B = \{4, 6\}$  and  $f = \{(1, 4), (5, 6), (8, 4), (9, 6)\}$ , then f is a surjective function.

a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.

c) A is true but R is false. d) A is false but R is true.

## Section B

21. For the principal value, evaluate  $\cot[\sin^{-1}\{\cos(\tan^{-1} 1)\}]$  [2]

OR

Which is greater,  $\tan 1$  or  $\tan^{-1} 1$ ?

22. Show that  $f(x) = \sin x - \cos x$  is an increasing function on  $(-\frac{\pi}{4}, \frac{\pi}{4})$ . [2]

23. A stone is dropped into a quiet lake and waves move in circles at a speed of 4 cm/s. At the instant, when the radius of the circular wave is 10 cm, how fast is the enclosed area increasing? [2]

OR

Show that the function  $f(x) = x^{100} + \sin x - 1$  is increasing on the interval  $(\frac{\pi}{2}, \pi)$

24. Evaluate:  $\int \tan^3 x \sec^3 x \, dx$  [2]

25. Prove that the determinant  $\begin{vmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{vmatrix}$  is independent of  $\theta$ . [2]

### Section C

26. Evaluate the integral:  $\int_0^\pi \frac{x}{a^2 \cos^2 x + b^2 \sin^2 x} dx$  [3]

27. In answering a question on a multiple choice questions test with four choices in each question, out of which only one is correct, a student either guesses or copies or knows the answer. The probability that he makes a guess is  $\frac{1}{4}$  and the probability he copies is also  $\frac{1}{4}$ . The probability that the answer is correct, given that he copied it is  $\frac{3}{4}$ . Find the probability that he knows the answer to the question, given that he correctly answered it. [3]

28. Evaluate the definite integral  $\int_0^{\frac{\pi}{4}} \frac{\sin x \cos x}{\cos^4 x + \sin^4 x} dx$  [3]

OR

$$\text{Evaluate the definite integral: } \int_1^2 e^{2x} \left( \frac{1}{x} - \frac{1}{2x^2} \right) dx$$

29. Solve the following differential equation  $\frac{dy}{dx} = 1 + x^2 + y^2 + x^2 y^2$ , given that  $y = 1$ , when  $x = 0$ . [3]

OR

Find the particular solution of the differential equation  $(xe^{x/y} + y)dx = x dy$ , given that  $y(1) = 0$

30. Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be three vectors such that  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$ ,  $|\vec{c}| = 5$  and each one of them being  $\perp$  to the sum of the other two, find  $|\vec{a} + \vec{b} + \vec{c}|$  [3]

OR

If  $\vec{a} = (\hat{i} - \hat{j}), \vec{b} = (3\hat{j} - \hat{k})$  and  $\vec{c} = (7\hat{i} - \hat{k})$ , find a vector  $\vec{d}$  which is perpendicular to both  $\vec{a}$  and  $\vec{b}$  and for which  $\vec{c} \cdot \vec{d} = 1$ .

31. Find  $\frac{dy}{dx}$  of the function  $(\cos x)^y = (\cos y)^x$ . [3]

### Section D

32. Find the area of the region  $\{(x, y) : x^2 + y^2 \leq 4, x + y \geq 2\}$ . [5]

33. Let  $A = \{1, 2, 3, \dots, 9\}$  and  $R$  be the relation in  $A \times A$  defined by  $(a, b) R (c, d)$  if  $a + d = b + c$  for  $(a, b), (c, d)$  in  $A \times A$ . Prove that  $R$  is an equivalence relation and also obtain the equivalence class  $[(2, 5)]$ . [5]

OR

Let  $A = R - \{3\}$  and  $B = R - \{1\}$ . Consider the function  $f: A \Rightarrow B$  defined by  $f(x) = \left( \frac{x-2}{x-3} \right)$ . Is  $f$  one-one and onto?

Justify your answer.

34. If  $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$  then show that  $A^2 - 5A + 7I = 0$  and hence find  $A^4$ . [5]

35. Prove that the volume of the largest cone that can be inscribed in a sphere of radius  $R$  is  $\frac{8}{27}$  of the volume of the sphere. [5]

OR

A window is in the form of a rectangle surmounted by a semicircular opening. The total perimeter of the window is 10 m. Find the dimensions of the window to admit maximum light through the whole opening.

### Section E

36. **Read the following text carefully and answer the questions that follow:** [4]

A shopkeeper sells three types of flower seeds  $A_1, A_2, A_3$ . They are sold in the form of a mixture, where the proportions of these seeds are  $4 : 4 : 2$  respectively. The germination rates of the three types of seeds are 45%, 60% and 35% respectively.



Based on the above information:


- Calculate the probability that a randomly chosen seed will germinate. (1)
- Calculate the probability that the seed is of type A2, given that a randomly chosen seed germinates. (1)
- A die is thrown and a card is selected at random from a deck of 52 playing cards. Then find the probability of getting an even number on the die and a spade card. (2)

**OR**

If A and B are any two events such that  $P(A) + P(B) - P(A \text{ and } B) = P(A|B)$ , then find  $P(A|B)$ . (2)

37. **Read the following text carefully and answer the questions that follow:** [4]

Consider the following diagram, where the forces in the cable are given.



- What is the equation of the line along cable AD? (1)
- What is length of cable DC? (1)
- Find vector DB (2)


**OR**

What is sum of vectors along the cable? (2)

38. **Read the following text carefully and answer the questions that follow:** [4]

Linear programming is a method for finding the optimal values (maximum or minimum) of quantities subject to the constraints when a relationship is expressed as linear equations or inequations.

- At which points is the optimal value of the objective function attained? (1)
- What does the graph of the inequality  $3x + 4y < 12$  look like? (1)
- Where does the maximum of the objective function  $Z = 2x + 5y$  occur in relation to the feasible region shown in the figure for the given LPP? (2)



**OR**

What are the conditions on the positive values of  $p$  and  $q$  that ensure the maximum of the objective function  $Z = px + qy$  occurs at both the corner points  $(15, 15)$  and  $(0, 20)$  of the feasible region determined by the given system of linear constraints? (2)

# Solution

## Section A

1.

(c) 
$$\begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a^n \end{bmatrix}$$

**Explanation:**  $A = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$

$$A^n = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix} \times \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix} \times \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix} \times \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix} \dots \{n \text{ times, (where } n \in \mathbb{N}\}$$

$$A^n = \begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a^n \end{bmatrix}$$

2. (a) 1

**Explanation:** When a given matrix is singular then the given matrix determinant is 0.

$$|A| = 0$$

$$\text{Given, } A = \begin{pmatrix} 3 - 2x & x + 1 \\ 2 & 4 \end{pmatrix}$$

$$|A| = 0$$

$$4(3 - 2x) - 2(x + 1) = 0$$

$$12 - 8x - 2x - 2 = 0$$

$$10 - 10x = 0$$

$$10(1 - x) = 0$$

$$x = 1$$

3.

(b)  $(A + B)^{-1} = B^{-1} + A^{-1}$

**Explanation:** Since, A and B are invertible matrices.

So, we can say that

$$(AB)^{-1} = B^{-1} A^{-1} \dots (i)$$

$$\text{We know that, } A^{-1} = \frac{1}{|A|} (\text{adj } A)$$

$$\Rightarrow \text{adj } A = |A| \cdot A^{-1} \dots (ii)$$

$$\text{Also, } \det(A)^{-1} = [\det(A)]^{-1}$$

$$\Rightarrow \det(A)^{-1} = \frac{1}{[\det(A)]}$$

$$\Rightarrow \det(A) \cdot \det(A)^{-1} = 1 \dots (iii)$$

Which is true,

So, only option d is incorrect.

4.

(c)  $f(x)$  is discontinuous  $\forall x \in \mathbb{Z} - \{1\}$

**Explanation:**  $f(x)$  is discontinuous  $\forall x \in \mathbb{Z} - \{1\}$

5. (a)  $\vec{r} = \hat{i} + 2\hat{j} + 3\hat{k} + \lambda(3\hat{i} + 2\hat{j} - 2\hat{k}), \lambda \in \mathbb{R}$

**Explanation:** The equation of the line which passes through the point  $(1, 2, 3)$  and is parallel to the vector  $3\hat{i} + 2\hat{j} - 2\hat{k}$ , let vector  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and vector  $\vec{b} = 3\hat{i} + 2\hat{j} - 2\hat{k}$ ,

the equation of line is :


$$\vec{a} + \lambda \vec{b} = (\hat{i} + \hat{j} + \hat{k}) + \lambda(3\hat{i} + 2\hat{j} - 2\hat{k})$$

6.

(c) 2

**Explanation:** Let the equation of given family be  $(x - h)^2 + (y - k)^2 = a^2$ . It has two arbitrary constants h and k. Therefore, the order of the given differential equation will be 2.

7. (a) at infinite number of points



**Explanation:**

Feasible region is shaded region with corner points  $(0, 0)$ ,  $(2, 0)$  and  $(0, 2)$

$$Z(0, 0) = 0$$

$$Z(2, 0) = 2 \leftarrow \text{maximise}$$

$$Z(0, 2) = 2 \leftarrow \text{maximise}$$

$Z_{\max} = 2$  obtained at  $(2, 0)$  and  $(0, 2)$  so is obtained at any point on line segment joining  $(2, 0)$  and  $(0, 2)$ .

8.

(c)  $[0, 1]$

**Explanation:** We have  $f(x) = \cos^{-1}(2x - 1)$

$$\text{Since, } -1 \leq 2x - 1 \leq 1$$

$$\Rightarrow 0 \leq 2x \leq 2$$

$$\Rightarrow 0 \leq x \leq 1$$

$$\therefore x \in [0, 1]$$

9.

(c)  $\log\left(\frac{4}{3}\right)$

**Explanation:**  $\log\left(\frac{4}{3}\right)$

$$\text{Let } I = \int_0^{\frac{\pi}{2}} \frac{\cos x}{(2+\sin x)(1+\sin x)} dx$$

Let  $\sin x = t$ , then  $\cos x dx = dt$

$$\text{When } x = 0, t = 0 \quad x = \frac{\pi}{2}, t = 1$$

Therefore the integral becomes

$$\begin{aligned} I &= \int_0^1 \frac{dt}{(2+t)(1+t)} \\ &= \int_0^1 \left[ \frac{-1}{2+t} + \frac{1}{1+t} \right] dt \\ &= [-\log(2+t) + \log(1+t)]_0^1 \\ &= [\log(1+t) - \log(2+t)]_0^1 \\ &= \log 2 - \log 3 - \log 1 + \log 2 \\ &= \log \frac{4}{3} \end{aligned}$$

10.

(c)  $\pm 4\sqrt{3}$

$$\text{Explanation: Given, } \begin{bmatrix} x & -5 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ 1 \end{bmatrix} = 0$$

$$\Rightarrow x \times 1 + (-5) \times 0 + (-1) \times 2 x \times 0 + (-5) \times 2 + (-1) \times 0$$

$$x \times 2 + (-5) \times 1 + (-1) \times 3 \begin{bmatrix} x \\ 4 \\ 1 \end{bmatrix} = 0$$

$$\Rightarrow [x - 2 \quad -10 \quad 2x - 8] \begin{bmatrix} x \\ 4 \\ 1 \end{bmatrix} = 0$$

$$\Rightarrow [(x - 2) \times x + (-10) \times 4 + (2x - 8) \times 1] = 0$$

$$\Rightarrow x^2 - 2x - 40 + 2x - 8 = 0$$

$$\Rightarrow x^2 = 48$$

$$\Rightarrow x = \pm\sqrt{48} = \pm 4\sqrt{3}$$

11. (a) a function to be optimized

**Explanation:** a function to be optimized

The objective function of a linear programming problem is either to be maximized or minimized i.e. objective function is to be optimized.

12.

(b)  $3(\hat{i} - 2\hat{j} + 2\hat{k})$

**Explanation:** Let  $\vec{a} = \hat{i} - 2\hat{j} + 2\hat{k}$

Unit vector in the direction of a vector  $\vec{a}$

$$= \frac{\vec{a}}{|\vec{a}|} = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{1^2 + 2^2 + 2^2}} = \frac{i - 2j + 2k}{3}$$

$\therefore$  Vector in the direction of  $\vec{a}$  with magnitude 9

$$= 9 \cdot \frac{i - 2j + 2k}{3} = 3(i - 2j + 2k).$$

13. (a) 14641

**Explanation:** We know that, for a square matrix of order n, if  $|A| \neq 0$

$$\text{Adj}(\text{Adj } A) = |A|^{n-2} A \ (\because n = 3)$$

$$\therefore \text{Adj}(\text{Adj } A) = |A|^{3-2} A \ (\because n = 3)$$

$$= |A| A$$

$$\therefore |\text{Adj}(\text{Adj } A)| = |A| |A| = |A|^3 \det A |A|^4$$

$$= 11^4 = 14641$$

14. (a)  $\frac{3}{8}$

**Explanation:** Given that,

$$P(A) = \frac{1}{5}, P(A \cup B) = \frac{7}{10}$$

Also, A and B are independent events,

$$\therefore P(A \cap B) = P(A) \cdot P(B)$$

$$\Rightarrow P(A) + P(B) - P(A \cup B) = P(A) \cdot P(B)$$

$$\Rightarrow \frac{1}{5} + P(B) - \frac{7}{10} = \frac{1}{5} \times P(B)$$

$$\Rightarrow P(B) - \frac{P(B)}{5} = \frac{7}{10} - \frac{1}{5} = \frac{5}{10}$$

$$\Rightarrow \frac{4P(B)}{5} = \frac{1}{2} \Rightarrow P(B) = \frac{5}{8}$$

$$\therefore P(\bar{B}) = 1 - P(B) = 1 - \frac{5}{8} = \frac{3}{8}$$

15.

(b) not defined

**Explanation:** not defined

16.

(b) 20

**Explanation:** We know that

$$(\vec{a} \cdot \vec{b})^2 + (\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$$

$$|\vec{a}|^2 \cdot |\vec{b}|^2 = 2^2 + 4^2$$

$$|\vec{a}|^2 \cdot |\vec{b}|^2 = 20$$

17.

(d)  $\frac{-1}{2}$ **Explanation:** Given that  $y = \tan^{-1} \left( \frac{\cos x}{1+\sin x} \right)$ Using  $\cos x = \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}$ ,  $\sin x = 2 \sin \frac{x}{2} \cos \frac{x}{2}$  and  $\cos^2 \theta + \sin^2 \theta = 1$ 

Therefore,

$$y = \tan^{-1} \left( \frac{\cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2} + 2 \sin \frac{x}{2} \cos \frac{x}{2}} \right) = \tan^{-1} \left( \frac{(\cos \frac{x}{2} - \sin \frac{x}{2})(\cos \frac{x}{2} + \sin \frac{x}{2})}{(\cos \frac{x}{2} + \sin \frac{x}{2})^2} \right)$$

$$\Rightarrow y = \tan^{-1} \frac{\cos \frac{x}{2} - \sin \frac{x}{2}}{\cos \frac{x}{2} + \sin \frac{x}{2}}$$

Dividing by  $\cos \frac{x}{2}$  in numerator and denominator, we get

$$y = \tan^{-1} \frac{1 - \tan \frac{x}{2}}{1 + \tan \frac{x}{2}}$$

Using  $\tan \left( \frac{\pi}{4} - x \right) = \frac{1 - \tan x}{1 + \tan x}$ , we obtain

$$y = \tan^{-1} \tan \left( \frac{\pi}{4} - \frac{x}{2} \right)$$

$$= \frac{\pi}{4} - \frac{x}{2}$$

Differentiating with respect to x, we

$$\frac{dy}{dx} = -\frac{1}{2}$$

18.

(c)  $\frac{\sqrt{3}}{\sqrt{55}}, \frac{4}{\sqrt{55}}, \frac{6}{\sqrt{55}}$ **Explanation:** Rewrite the given line as

$$r \frac{2(x - \frac{1}{2})}{\sqrt{3}} = \frac{y+2}{2} = \frac{z-3}{3}$$

$$\text{or } \frac{x - \frac{1}{2}}{\sqrt{3}} = \frac{y+2}{4} = \frac{z-3}{6}$$

 $\therefore$  DR's of line are  $\sqrt{3}, 4$  and  $6$ 

Therefore, direction cosines are:

$$\frac{\sqrt{3}}{\sqrt{(\sqrt{3})^2 + 4^2 + 6^2}}, \frac{4}{\sqrt{(\sqrt{3})^2 + 4^2 + 6^2}}, \frac{6}{\sqrt{(\sqrt{3})^2 + 4^2 + 6^2}} \text{ or } \frac{\sqrt{3}}{\sqrt{55}}, \frac{4}{\sqrt{55}}, \frac{6}{\sqrt{55}}$$

19.

(c) A is true but R is false.

**Explanation:** Let  $S(x)$  be the selling price of  $x$  items and let  $C(x)$  be the cost price of  $x$  items.

Then, we have

$$S(x) = (5 - \frac{x}{100})x = 5x - \frac{x^2}{100}$$

$$\text{and } C(x) = \frac{x}{5} + 500$$

Thus, the profit function  $P(x)$  is given by

$$P(x) = S(x) - C(x) = 5x - \frac{x^2}{100} - \frac{x}{5} - 500$$

$$\text{i.e. } P(x) = \frac{24}{5}x - \frac{x^2}{100} - 500$$

On differentiating both sides w.r.t.  $x$ , we get

$$P'(x) = \frac{24}{5} - \frac{x}{50}$$

Now,  $P'(x) = 0$  gives  $x = 240$ .

$$\text{Also, } P'(x) = \frac{-1}{50}.$$

$$\text{So, } P'(240) = \frac{-1}{50} < 0$$

Thus,  $x = 240$  is a point of maxima.

Hence, the manufacturer can earn maximum profit, if he sells 240 items.

20.

(d) A is false but R is true.

**Explanation:** We have,  $A = \{1, 5, 8, 9\}$ ,  $B = \{4, 6\}$  and  $f = \{(1, 4), (5, 6), (8, 4), (9, 6)\}$ So, all elements of  $B$  has a domain element on  $A$  or we can say elements 1 and 8 & 5 and 9 have some range 4 & 6, respectively.

Therefore,  $f : A \rightarrow B$  is a surjective function not one to one function.

Also, for a bijective function,  $f$  must be both one to one onto.

### Section B

21. We know that  $\tan^{-1} 1 = \frac{\pi}{4}$ .


$$\begin{aligned} \therefore \cot[\sin^{-1}\{\cos(\tan^{-1} 1)\}] \\ = \cot\left\{\sin^{-1}\left(\cos\frac{\pi}{4}\right)\right\} = \cot\left(\sin^{-1}\frac{1}{\sqrt{2}}\right) = \cot\frac{\pi}{4} = 1 \end{aligned}$$

OR

From Fig. we note that  $\tan x$  is an increasing function in the interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$ , since  $1 > \frac{\pi}{4} \Rightarrow \tan 1 > \tan \frac{\pi}{4}$ . This gives  $\tan 1 > 1$

$$\Rightarrow \tan 1 > 1 > \frac{\pi}{4}$$

$$\Rightarrow \tan 1 > 1 > \tan^{-1}(1)$$



22. Given:  $f(x) = \sin x - \cos x$

$$f(x) = \cos x + \sin x$$

$$= \sqrt{2} \left( \frac{1}{\sqrt{2}} \cos x + \frac{1}{\sqrt{2}} \sin x \right)$$

$$= \sqrt{2} \left( \frac{\sin \pi}{4} \cos x + \frac{\cos \pi}{4} \sin x \right)$$

$$= \sqrt{2} \sin\left(\frac{\pi}{4} + x\right)$$

Now,

$$x \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$$

$$\Rightarrow -\frac{\pi}{4} < x < \frac{\pi}{4}$$

$$\Rightarrow 0 < \frac{\pi}{4} + x < \frac{\pi}{2}$$

$$\Rightarrow \sin 0^\circ < \sin\left(\frac{\pi}{4} + x\right) < \sin \frac{\pi}{2}$$

$$\Rightarrow 0 < \sin\left(\frac{\pi}{4} + x\right) < 1$$

$$\Rightarrow \sqrt{2} \sin\left(\frac{\pi}{4} + x\right) > 0$$

$$\Rightarrow f(x) > 0$$

Hence,  $f(x)$  is an increasing function on  $(-\frac{\pi}{4}, \frac{\pi}{4})$

23. Let  $A$  be the area of the circle of radius  $r$ .

$$\text{Then, } A = \pi r^2$$

Therefore, the rate of change of area  $A$  with respect to time 't' is

$$\frac{dA}{dt} = \frac{d}{dt}(\pi r^2) = \frac{d}{dr}(\pi r^2) \cdot \frac{dr}{dt} = 2\pi r \frac{dr}{dt} \quad \text{...(By Chain Rule)}$$

$$\text{Given that } \frac{dr}{dt} = 4 \text{ cm/s}$$

$$\text{Therefore, when } r = 10, \frac{dA}{dt} = 2\pi \times 10 \times 4 = 80\pi$$

Thus, the enclosed area is increasing at a rate of  $80\pi \text{ cm}^2/\text{s}$ , when  $r = 10 \text{ cm}$ .

OR

Given interval:  $x \in (\pi/2, \pi)$

$$\Rightarrow \pi/2 < x < \pi$$

$$x^{99} > 1$$

$$100x^{99} > 100$$

Again,  $x \in (\pi/2, \pi) \Rightarrow -1 < \cos x < 0 \Rightarrow 0 > \cos x > -1$

$$100x^{99} > 100 \text{ and } \cos x > -1$$

$$100x^{99} + \cos x > 100 - 1 = 99$$

$$100x^{99} + \cos x > 0$$

$$f'(x) > 0$$

Thus  $f(x)$  is increasing on  $(\pi/2, \pi)$

24. Let  $I = \int \tan^3 x \sec^3 x \, dx$ , then we have

$$I = \int \tan^2 x \sec^2 x (\sec x \tan x) \, dx = \int (\sec^2 x - 1) \sec^2 x (\sec x \tan x) \, dx$$

Substituting  $\sec x = t$  and  $\sec x \tan x \, dx = dt$ , we obtain

$$I = \int (t^2 - 1) t^2 \, dt = \int (t^4 - t^2) \, dt = \frac{t^5}{5} - \frac{t^3}{3} + C = \frac{1}{5} \sec^5 x - \frac{1}{3} \sec^3 x + C$$

$$25. \text{ Let } \Delta = \begin{bmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{bmatrix}$$

Expanding along first row,

$$\Delta = x \begin{vmatrix} -x & 1 \\ 1 & x \end{vmatrix} - \sin \theta \begin{vmatrix} -\sin \theta & 1 \\ \cos \theta & x \end{vmatrix} + \cos \theta \begin{vmatrix} -\sin \theta & -x \\ \cos \theta & 1 \end{vmatrix}$$

$$\Rightarrow \Delta = x(-x^2 - 1) - \sin \theta(-x \sin \theta - \cos \theta) + \cos \theta(-\sin \theta + x \cos \theta)$$

$$\Rightarrow \Delta = -x^3 - x + x \sin^2 \theta + \sin \theta \cos \theta - \sin \theta \cos \theta + x \cos^2 \theta$$

$$\Rightarrow \Delta = -x^3 - x + x(\sin^2 \theta + \cos^2 \theta) = -x^3 - x + x = -x^3 \text{ which is independent of } \theta$$

### Section C

26. We have,

$$I = \int_0^\pi \frac{x}{a^2 \cos^2 x + b^2 \sin^2 x} \, dx \dots (i)$$

Using property of definite integral we have,

$$= \int_0^\pi \frac{(\pi - x)}{a^2 \cos^2(\pi - x) + b^2 \sin^2(\pi - x)} \, dx$$

$$= \int_0^\pi \frac{\pi - x}{a^2 \cos^2 x + b^2 \sin^2 x} \, dx \dots (ii)$$

Adding (i) and (ii)

$$2I = \int_0^\pi \frac{x + \pi - x}{a^2 \cos^2 x + b^2 \sin^2 x} \, dx$$

$$= \pi \int_0^\pi \frac{1}{a^2 \cos^2 x + b^2 \sin^2 x} \, dx$$

$$= \pi \int_0^\pi \frac{\sec^2 x}{a^2 + b^2 \tan^2 x} \, dx \dots (\text{Dividing numerator and denominator by } \cos^2 x)$$

$$= 2\pi \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{a^2 + b^2 \tan^2 x} \, dx \dots \left[ \text{Using } \int_0^{2a} f(x) \, dx = \int_0^a f(x) \, dx + \int_0^a f(2a - x) \, dx \right]$$

Put  $\tan x = t$

$$\Rightarrow \sec^2 x \, dx = dt$$

When  $x \rightarrow 0; t \rightarrow 0$

and  $x \rightarrow \frac{\pi}{2}; t \rightarrow \infty$

$$\therefore 2I = 2\pi \int_0^{\frac{\pi}{2}} \frac{dt}{a^2 + b^2 t^2}$$

$$\Rightarrow I = \frac{\pi}{b^2} \int_0^{\frac{\pi}{2}} \frac{dt}{\frac{a^2}{b^2} + t^2}$$

$$= \frac{\pi}{b^2} \times \frac{b}{a} \left[ \tan^{-1} \left( \frac{bt}{a} \right) \right]_0^\infty$$

$$= \frac{\pi}{ab} \left[ \frac{\pi}{2} - 0 \right]$$

$$= \frac{\pi}{ab} \times \frac{\pi}{2}$$

$$= \frac{\pi^2}{2ab}$$

$$\text{Hence, } I = \frac{\pi^2}{2ab}$$

27. Let  $E_1$  = Student guesses the answer

$E_2$  = Student copies the answer

$E_3$  = Student knows the answer

$A$  = Student answers the question correctly.

$$P(E_1) = \frac{1}{4}, P(E_2) = \frac{1}{4}, P(E_3) = 1 - \left( \frac{1}{4} + \frac{1}{4} \right) = \frac{1}{2}$$

$$P(A | E_1) = \frac{1}{4}, P(A | E_2) = \frac{3}{4}, P(A | E_3) = 1$$

The required probability

$$= P(E_3 | A) = \frac{P(E_3) \times P(A | E_3)}{\sum_{i=1}^3 P(E_i) \times P(A | E_i)}$$

$$\begin{aligned}
&= \frac{\frac{1}{2} \times 1}{\frac{1}{4} \times \frac{1}{4} + \frac{1}{4} \times \frac{3}{4} + \frac{1}{2} \times 1} \\
&= \frac{1}{\frac{1}{8} + \frac{3}{8} + 1} = \frac{8}{12} = \frac{2}{3}
\end{aligned}$$

28.  $I = \int_0^{\pi/4} \frac{\sin x \cos x}{\cos^4 x + \sin^4 x} dx$

Dividing Nr. and Dr. by  $\cos^4 x$

$$\begin{aligned}
&= \int_0^{\pi/4} \frac{\frac{\sin x \cos x}{\cos^4 x}}{\frac{\cos^4 x}{\cos^4 x} + \frac{\sin^4 x}{\cos^4 x}} dx \\
&= \int_0^{\pi/4} \frac{\tan x \sec^2 x}{1 + \tan^4 x} dx \\
&= \int_0^{\pi/4} \frac{\tan x \sec^2 x}{1 + (\tan^2 x)^2} dx
\end{aligned}$$

Put  $\tan^2 x = t$

$$2 \tan x \sec^2 x dx = dt$$

When  $x = 0, t = 0$  and when  $x = \frac{\pi}{4}, t = 1$

$$\begin{aligned}
\therefore I &= \frac{1}{2} \int_0^1 \frac{dt}{1+t^2} \\
&= \frac{1}{2} [\tan^{-1} t]_0^1 \\
&= \frac{1}{2} \cdot \frac{\pi}{4} = \frac{\pi}{8}
\end{aligned}$$

OR

We have,

$$\begin{aligned}
I &= \int_1^2 e^{2x} \left( \frac{1}{x} - \frac{1}{2x^2} \right) dx \\
I &= \int_1^2 \frac{1}{x} \cdot e^{2x} - \int_1^2 \frac{1}{2x^2} \cdot e^{2x} dx \\
\Rightarrow I &= I_1 - I_2
\end{aligned}$$

Now,  $I_1 = \int_1^2 \frac{1}{x} e^{2x}$  (By parts we have)

$$\begin{aligned}
\Rightarrow I_1 &= \left[ \frac{1}{x} \right]_1^2 \cdot \int_1^2 e^{2x} dx - \int_1^2 -\frac{1}{x^2} \frac{e^{2x}}{2} dx \\
\Rightarrow I_1 &= \left[ \frac{1}{x} \cdot \frac{e^{2x}}{2} \right]_1^2 + \int_1^2 \frac{1}{2x^2} e^{2x} dx \\
\Rightarrow I_1 &= \left[ \frac{1}{2x} e^{2x} \right]_1^2 + I_2
\end{aligned}$$

As,  $I = I_1 - I_2$

$$\begin{aligned}
\Rightarrow I &= \left[ \frac{1}{2x} e^{2x} \right]_1^2 - I_2 + I_2 \\
\Rightarrow I &= \left[ \frac{1}{2x} e^{2x} \right]_1^2 = \frac{1}{2} \left[ \frac{1}{2} e^4 - e^2 \right] \\
\Rightarrow I &= \frac{1}{4} e^2 (e^2 - 1)
\end{aligned}$$

29. According to the question ,

Given differential equation is ,

$$\begin{aligned}
\frac{dy}{dx} &= 1 + x^2 + y^2 + x^2 y^2 \\
\Rightarrow \frac{dy}{dx} &= 1 (1 + x^2) + y^2 (1 + x^2) \\
\Rightarrow \frac{dy}{dx} &= (1 + x^2) (1 + y^2) \\
\Rightarrow \frac{dy}{1+y^2} &= (1 + x^2) dx
\end{aligned}$$

On integrating both sides, we get

$$\begin{aligned}
\int \frac{dy}{1+y^2} &= \int (1 + x^2) dx \\
\Rightarrow \tan^{-1} y &= x + \frac{x^3}{3} + C \quad \dots(i)
\end{aligned}$$

Given that  $y = 1$ , when  $x = 0$ .

On putting  $x = 0$  and  $y = 1$  in Eq. (i), we get

$$\begin{aligned}
\tan^{-1} 1 &= C \\
\Rightarrow \tan^{-1} (\tan \pi/4) &= C \quad \left[ \because \tan \frac{\pi}{4} = 1 \right] \\
\Rightarrow C &= \frac{\pi}{4}
\end{aligned}$$

On putting the value of  $C$  in Eq. (i), we get

$$\tan^{-1} y = x + \frac{x^3}{3} + \frac{\pi}{4}$$

$$\therefore y = \tan\left(x + \frac{x^3}{3} + \frac{\pi}{4}\right)$$

which is the required solution of differential equation.

OR

The given differential equation can be rewritten as,

$$\begin{aligned} xe^{\frac{y}{x}} - y + x \frac{dy}{dx} &= 0 \\ \Rightarrow x \frac{dy}{dx} &= y - xe^{\frac{y}{x}} \\ \Rightarrow \frac{dy}{dx} &= \left(\frac{y}{x}\right) - e^{\frac{y}{x}} \\ \Rightarrow \frac{dy}{dx} &= f\left(\frac{y}{x}\right) \end{aligned}$$

$\Rightarrow$  the given differential equation is a homogenous equation.

The solution of the given differential equation is:

Put  $y = vx$

$$\begin{aligned} \Rightarrow \frac{dy}{dx} &= v + x \frac{dv}{dx} \\ v + x \frac{dv}{dx} &= \left(\frac{vx}{x}\right) - e^{\frac{vx}{x}} \\ \Rightarrow x \frac{dv}{dx} &= -e^v \\ \Rightarrow \frac{dv}{e^v} &= \frac{-dx}{x} \end{aligned}$$

Integrating both the sides we get:

$$\begin{aligned} \Rightarrow \int \frac{dv}{e^v} &= - \int \frac{dx}{x} + c \\ \Rightarrow -e^{-v} &= -\ln|x| + c \end{aligned}$$

Resubstituting the value of  $y = vx$  we get

$$\Rightarrow -e^{-\left(\frac{y}{x}\right)} = -\ln|x| + c$$

Now,  $y(1) = 0$

$$\Rightarrow -e^{-(0)} = -\ln|1| + c$$

$$\Rightarrow c = -1$$

$$\Rightarrow \log|x| + e^{-y/x} = 1$$

$$30. \vec{a} \cdot (\vec{b} + \vec{c}) = 0, \vec{b} \cdot (\vec{c} + \vec{a}) = 0, \vec{c} \cdot (\vec{a} + \vec{b}) = 0 \text{ (Given)}$$

$$\begin{aligned} |\vec{a} + \vec{b} + \vec{c}|^2 &= (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) \\ &= \vec{a} \cdot \vec{a} + \vec{a} \cdot (\vec{b} + \vec{c}) + \vec{b} \cdot \vec{b} + \vec{b} \cdot (\vec{a} + \vec{c}) + \vec{c} \cdot \vec{c} + \vec{c} \cdot (\vec{a} + \vec{b}) \\ &= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 \end{aligned}$$

$$= 9 + 16 + 25$$

$$= 50$$

$$|\vec{a} + \vec{b} + \vec{c}| = \sqrt{50}$$

$$= 5\sqrt{2}$$

OR

$$\text{Let } \vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$$

$$\vec{d} \perp \vec{a}, \vec{d} \cdot \vec{a} = 0 \Rightarrow (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) \cdot (\hat{i} - \hat{j}) = 0$$

$$\Rightarrow a_1 - a_2 = 0 \dots (i)$$

$$\vec{d} \perp \vec{b}, \vec{a} \cdot \vec{b} = 0 \Rightarrow (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) \cdot (3\hat{j} - \hat{k}) = 0$$

$$\Rightarrow 3a_2 - a_3 = 0 \dots (ii)$$

$$\vec{d} \cdot \vec{c} = 1 \Rightarrow (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) \cdot (7\hat{i} - \hat{k}) = 1$$

$$\Rightarrow 7a_1 - a_3 = 1 \dots (iii)$$

Solving equation (i) and (ii) we get  $3a_1 - a_3 = 0 \dots (iv)$

Again solving equation (iii) & (iv) we get  $a_1 = \frac{1}{4}$

From equation (i),  $a_1 - a_2 = 0$  or  $a_1 = a_2 = \frac{1}{4}$

From equation (ii),  $3a_2 - a_3 = 0 \Rightarrow 3 \cdot \frac{1}{4} = a_3 \Rightarrow a_3 = \frac{3}{4}$

Hence,  $\vec{d} = \frac{1}{4} \hat{i} + \frac{1}{4} \hat{j} + \frac{3}{4} \hat{k}$

31. We have,  $(\cos x)^y = (\cos y)^x$



Which is true for any  $a, b \in A$

Hence,  $R$  is reflexive.

Let  $(a, b) R (c, d)$

$$a+d = b+c$$

$$c+b = d+a \Rightarrow (c, d) R (a, b)$$

So,  $R$  is symmetric.

Let  $(a, b) R (c, d)$  and  $(c, d) R (e, f)$

$$a+d = b+c \text{ and } c+f = d+e$$

$$a+d = b+c \text{ and } d+e = c+f \Rightarrow (a+d) - (d+e) = (b+c) - (c+f)$$

$$(a-e) = b-f$$

$$a+f = b+e$$

$$(a, b) R (e, f)$$

So,  $R$  is transitive.

Hence  $R$  is an equivalence relation.

Let  $(a, b) R (2, 5)$ , then

$$a+5 = b+2$$

$$a = b-3$$

If  $b < 3$ , then  $a$  does not belong to  $A$ .

Therefore, possible values of  $b$  are  $> 3$ .

For  $b = 4, 5, 6, 7, 8, 9$

$$a = 1, 2, 3, 4, 5, 6$$

Therefore, equivalence class of  $(2, 5)$  is

$$\{(1, 4), (2, 5), (3, 6), (4, 7), (5, 8), (6, 9)\}$$

OR

$$A = R - \{3\} \text{ and } B = R - \{1\} \text{ and } f(x) = \left( \frac{x-2}{x-3} \right)$$

$$\text{Let } x_1, x_2 \in A, \text{ then } f(x_1) = \frac{x_1-2}{x_1-3} \text{ and } f(x_2) = \frac{x_2-2}{x_2-3}$$

Now, for  $f(x_1) = f(x_2)$

$$\Rightarrow \frac{x_1-2}{x_1-3} = \frac{x_2-3}{x_2-3}$$

$$\Rightarrow (x_1-2)(x_2-3) = (x_2-2)(x_1-3)$$

$$\Rightarrow x_1x_2 - 3x_1 - 2x_2 + 6 = x_1x_2 - 2x_1 - 3x_2 + 6$$

$$\Rightarrow -3x_1 - 2x_2 = -2x_1 - 3x_2$$

$$= x_1 = x_2$$

$\therefore f$  is one-one function.

$$\text{Now } y = \frac{x-2}{x-3}$$

$$\Rightarrow y(x-3) = x-2$$

$$\Rightarrow xy - 3y = x - 2$$

$$\Rightarrow x(y-1) = 3y - 2$$

$$\Rightarrow x = \frac{3y-2}{y-1}$$

$$\therefore f\left(\frac{3y-2}{y-1}\right) = \frac{\frac{3y-2}{y-1}-2}{\frac{3y-2}{y-1}-3} = \frac{3y-2-2y+2}{2y-2-3y+3} = y$$

$$\Rightarrow f(x) = y$$

Therefore,  $f$  is an onto function.

34. Given  $A^2 - 5A + 7I = 0$

$$\text{L.H.S} = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \text{R.H.S}$$

$$A^2 = 5A - 7I$$

$$A^3 = A^2 \cdot A$$

$$= 5A^2 - 7AI$$

$$= 5A^2 - 7A \text{ (Since } AI = A)$$

$$= 5(5A - 7I) - 7A$$

$$= 25A - 35I - 7A$$

$$A^3 = 18A - 35I$$

$$A^4 = A^3 \cdot A$$

$$= (18A - 35I) \cdot A$$

$$= 18A^2 - 35IA$$

$$= 18(5A - 7I) - 35A$$


$$= 90A - 126I - 35A$$

$$= 55A - 126I$$

$$= 55 \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} - 126 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 165 & 55 \\ -55 & 110 \end{bmatrix} + \begin{bmatrix} -126 & 0 \\ 0 & -126 \end{bmatrix}$$

$$A^4 = \begin{bmatrix} 39 & 55 \\ -55 & -16 \end{bmatrix}.$$



$$v = \frac{1}{3}\pi r^2 h \quad [r^2 = \sqrt{R^2 - x^2}]$$

$$V = \frac{1}{2}\pi \cdot (R^2 - x^2) \cdot (R + x)$$

$$\frac{dy}{dx} = \frac{1}{3}\pi [(R^2 - x^2)(1) + (R + x)(-2x)]$$

$$= \frac{1}{3}\pi [(R + x)(R - x) - 2x(R + x)]$$

$$= \frac{1}{3}\pi (R + x)[R - x - 2x]$$

$$= \frac{1}{3}\pi (R + x)(R - 3x) \dots (1)$$

$$\text{Put } \frac{dv}{dr} = 0$$

$$R = -x \text{ (neglecting)}$$

$$R = 3x$$

$$\frac{R}{3} = x$$

On again differentiating equation (1)

$$\frac{d^2v}{dx^2} = \frac{1}{3}\pi [(R + x)(-3) + (R - 3x)(1)]$$

$$= \frac{d^2v}{dx^2} \Big|_{x=\frac{R}{3}} = \frac{1}{3}\pi \left[ \left( R + \frac{R}{3} \right) (-3) + \left( R - 3 \cdot \frac{R}{3} \right) \right]$$

$$= \frac{1}{3}\pi \left[ \frac{4R}{3} \times -3 + 0 \right]$$

$$= \frac{-1}{3}\pi 4R$$

$$\frac{d^2v}{dx^2} < 0 \text{ Hence maximum}$$

$$\text{Now } v = \frac{1}{3}\pi [(R^2 - x^2)(R + x)] \quad [x = \frac{R}{3}]$$

$$v = \frac{1}{3}\pi \left[ \left( R^2 - \left( \frac{R}{3} \right)^2 \right) \left( R + \left( \frac{R}{3} \right) \right) \right]$$


$$= \frac{1}{3}\pi \left[ \frac{8R^2}{9} \times \frac{4R}{3} \right]$$

$$v = \frac{8}{27} \left( \frac{4}{3} \right) \pi R^3$$

$$v = \frac{8}{27} \text{ Volume of sphere}$$

$$\text{Volume of cone} = \frac{8}{27} \text{ of volume of sphere.}$$

OR



Let  $P$  be the perimeter of window

$$P = 2x + 2r + \frac{1}{2} \times 2\pi r$$

$$10 = 2x + 2r + \pi r \quad [P = 10]$$

$$x = \frac{10 - 2r - \pi r}{2}$$

Let  $A$  be area of window

$$A = 2rx + \frac{1}{2}\pi r^2$$

$$= 2r \left[ \frac{10 - 2r - \pi r}{2} \right] + \frac{1}{2}\pi r^2$$

$$= 10r - 2r^2 - \pi r^2 + \frac{1}{2}\pi r^2$$

$$= 10r - 2r^2 - \frac{\pi r^2}{2}$$

$$\frac{dA}{dr} = 10 - 4r - \pi r$$

$$\frac{d^2A}{dr^2} = -(\pi + 4)$$

$$\frac{dA}{dr} = 0$$

$$r = \frac{10}{\pi + 4}$$

$$\frac{d^2A}{dr^2} < 0 \text{ maximum}$$


$$x = \frac{10 - 2r - \pi r}{2}$$

$$x = \frac{10}{\pi + 4}$$

$$\text{Length of rectangle} = 2r = \frac{20}{\pi + 4}$$

$$\text{width} = \frac{10}{\pi + 4}$$

## Section E



$$\text{Here, } P(E_1) = \frac{4}{10}, P(E_2) = \frac{4}{10}, P(E_3) = \frac{2}{10}$$

$$P\left(\frac{A}{E_1}\right) = \frac{45}{100}, P\left(\frac{A}{E_2}\right) = \frac{60}{100}, P\left(\frac{A}{E_3}\right) = \frac{35}{100}$$

$$\therefore P(A) = P(E_1) \cdot P\left(\frac{A}{E_1}\right) + P(E_2) \cdot P\left(\frac{A}{E_2}\right) + P(E_3) \cdot P\left(\frac{A}{E_3}\right)$$

$$= \frac{4}{10} \times \frac{45}{100} + \frac{4}{10} \times \frac{60}{100} + \frac{2}{10} \times \frac{35}{100}$$

$$= \frac{180}{1000} + \frac{240}{1000} + \frac{70}{1000}$$

$$= \frac{490}{1000} = 4.9$$

$$\text{ii. Required probability} = P\left(\frac{E_2}{A}\right)$$

$$= \frac{P(E_2) \cdot P\left(\frac{A}{E_2}\right)}{P(A)}$$

$$= \frac{\frac{4}{10} \times \frac{60}{100}}{\frac{490}{1000}}$$

$$= \frac{240}{490} = \frac{24}{49}$$

iii. Let,

$E_1$  = Event for getting an even number on die and

$E_2$  = Event that a spade card is selected

$$\therefore P(E_1) = \frac{3}{6}$$

$$= \frac{1}{2}$$

$$\text{and } P(E_2) = \frac{13}{52} = \frac{1}{4}$$

Then,  $P(E_1 \cap E_2) = P(E_1) \cdot P(E_2)$

$$= \frac{1}{2}, \frac{1}{4} = \frac{1}{8}$$

**OR**

$$P(A) + P(B) - P(A \text{ and } B) = P(A)$$

$$\Rightarrow P(A) + P(B) - P(A \cap B) = P(A)$$

$$\Rightarrow P(B) - P(A \cap B) = 0$$

$$\Rightarrow P(A \cap B) = P(B)$$

$$\therefore P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$= \frac{P(B)}{P(B)}$$

$$= 1$$

37. i. Clearly, the coordinates of A are (8, 10, 0) and D are (0, 0, 30)

$\therefore$  Equation of AD is given by

$$\frac{x-0}{8-0} = \frac{y-0}{10-0} = \frac{z-30}{-30}$$

$$\Rightarrow \frac{x}{4} = \frac{y}{5} = \frac{30-z}{15}$$

ii. The coordinates of point C are (15, -20, 0) and D are (0, 0, 30)

$\therefore$  Length of the cable DC

$$= \sqrt{(0-15)^2 + (0+20)^2 + (30-0)^2}$$

$$= \sqrt{225 + 400 + 900} = \sqrt{1525} = 5\sqrt{61} \text{ m}$$

iii. Since, the coordinates of point B are (-6, 4, 0) and D are (0, 0, 30), therefore vector DB is

$$(-6-0)\hat{i} + (4-0)\hat{j} + (0-30)\hat{k}, \text{ i.e., } -6\hat{i} + 4\hat{j} - 30\hat{k}$$

**OR**


Required sum

$$= (8\hat{i} + 10\hat{j} - 30\hat{k}) + (-6\hat{i} + 4\hat{j} - 30\hat{k}) + (15\hat{i} - 20\hat{j} - 30\hat{k})$$

$$= 17\hat{i} - 6\hat{j} - 90\hat{k}$$

38. i. When we solve an L.P.P. graphically, the optimal (or optimum) value of the objective function is attained at corner points of the feasible region.

ii. From the graph of  $3x + 4y < 12$  it is clear that it contains the origin but not the points on the line  $3x + 4y = 12$ .



iii. Maximum of objective function occurs at corner points.

Corner Points	Value of $Z = 2x + 5y$
(0, 0)	0
(7, 0)	14
(6, 3)	27
(4, 5)	33 $\leftarrow$ Maximum
(0, 6)	30

**OR**

Value of  $Z = px + qy$  at (15, 15) =  $15p + 15q$  and that at (0, 20) =  $20q$ . According to given condition, we have

$$15p + 15q = 20q \Rightarrow 15p = 5q \Rightarrow q = 3p$$